

Cohomologies on Symplectic Manifolds

Raquel Villacampa

Joint work with Luis Ugarte (arXiv:1404.6777(math.SG))



Parma, November 28th 2014

PRINCIPAL IDEAS

- 1 There exist several cohomologies on symplectic manifolds: **harmonic, primitive, filtered and coeffective**. All of them appear independently in the literature.

Objective 1 Relate all these cohomologies by defining the **generalized coeffective cohomology**.

PRINCIPAL IDEAS

- 1 There exist several cohomologies on symplectic manifolds: **harmonic, primitive, filtered and coeffective**. All of them appear independently in the literature.

Objective 1 Relate all these cohomologies by defining the **generalized coeffective cohomology**.

- 2 Harmonic, primitive and filtered cohomologies are defined in terms of d^\wedge , ∂_+ , ∂_- and therefore these cohomologies are difficult to compute.

Objective 2 Provide a **coeffective version** of these cohomologies so that computations are easier.

PRINCIPAL IDEAS

- 1 There exist several cohomologies on symplectic manifolds: **harmonic, primitive, filtered and coeffective**. All of them appear independently in the literature.

Objective 1 Relate all these cohomologies by defining the **generalized coeffective cohomology**.

- 2 Harmonic, primitive and filtered cohomologies are defined in terms of d^\wedge , ∂_+ , ∂_- and therefore these cohomologies are difficult to compute.

Objective 2 Provide a **coeffective version** of these cohomologies so that computations are easier.

- 3 The dimension of all these cohomology groups can vary when the symplectic form does (notion of *flexibility*).

Objective 3 Study harmonic and filtered flexibility in terms of **coeffective flexibility**.

SYMPLECTIC HARMONICITY I

- (M^{2n}, ω) symplectic manifold.
- $\Omega^*(M)$ the space of differential forms on M .

[Brylinski 88] $\alpha \in \Omega^*(M)$ is **symplectically harmonic** if $d\alpha = 0 = d^\wedge \alpha$.
(d^\wedge is the adjoint of d with respect to $*_\omega$.)

- Notation: $\Omega_{\text{hr}}^*(M)$.

SYMPLECTIC HARMONICITY I

- (M^{2n}, ω) symplectic manifold.
- $\Omega^*(M)$ the space of differential forms on M .

[Brylinski 88] $\alpha \in \Omega^*(M)$ is **symplectically harmonic** if $d\alpha = 0 = d^\wedge \alpha$.
(d^\wedge is the adjoint of d with respect to $*_\omega$.)

- Notation: $\Omega_{\text{hr}}^*(M)$.
- **Problem**: there are non-zero exact symplectically harmonic forms (important difference w.r.t. the usual Hodge Theory).

SYMPLECTIC HARMONICITY I

- (M^{2n}, ω) symplectic manifold.
- $\Omega^*(M)$ the space of differential forms on M .

[Brylinski 88] $\alpha \in \Omega^*(M)$ is **symplectically harmonic** if $d\alpha = 0 = d^\wedge \alpha$.
(d^\wedge is the adjoint of d with respect to $*_\omega$.)

- Notation: $\Omega_{\text{hr}}^*(M)$.
- **Problem**: there are non-zero exact symplectically harmonic forms (important difference w.r.t. the usual Hodge Theory).

Symplectically harmonic cohomology: $H_{\text{hr}}^q(M) = \frac{\Omega_{\text{hr}}^q(M)}{(\Omega_{\text{hr}}^q(M) \cap \text{im } d)}$.

- $H_{\text{hr}}^q(M) \stackrel{?}{=} H^q(M)$, $q = 0, \dots, 2n$.

Yes \iff HLC ($L^k : H^{n-k} \xrightarrow{\cong} H^{n+k}$, $k = 1, \dots, n$).

[Mathieu 95], [Yan 96]

SYMPLECTIC HARMONICITY II

Theorem [Ibáñez-Rudyak-Tralle-Ugarte 01], [Yamada 02]

- $H_{\text{hr}}^q(M) = P^q(M, \omega) + L(H_{\text{hr}}^{q-2}(M)), \quad q = 0, \dots, n,$
- $H_{\text{hr}}^q(M) = \text{Im} \{L^{q-n} : H_{\text{hr}}^{2n-q}(M) \rightarrow H^q(M)\}, \quad q = n+1, \dots, 2n,$

where

$$P^q(M) = \{[\alpha] \in H^q(M) \mid L^{n-q+1}[\alpha] = 0\} \subset H_{\text{hr}}^q(M).$$

SYMPLECTIC HARMONICITY II

Theorem [Ibáñez-Rudyak-Tralle-Ugarte 01], [Yamada 02]

- $H_{\text{hr}}^q(M) = P^q(M, \omega) + L(H_{\text{hr}}^{q-2}(M)), \quad q = 0, \dots, n,$
- $H_{\text{hr}}^q(M) = \text{Im} \{L^{q-n} : H_{\text{hr}}^{2n-q}(M) \rightarrow H^q(M)\}, \quad q = n+1, \dots, 2n,$

where

$$P^q(M) = \{[\alpha] \in H^q(M) \mid L^{n-q+1}[\alpha] = 0\} \subset H_{\text{hr}}^q(M).$$

If (M^{2n}, ω) is of finite type: $h_q(M) = \dim H_{\text{hr}}^q(M)$ is finite.

- $h_i = b_i$ for $i = 0, 1, 2.$
- If (M^{2n}, ω) is closed: $h_{2n} = b_{2n}$, h_{2n-1} is even.

PRIMITIVE COHOMOLOGIES [Tseng-Yau I & II, 12]

New finite dimensional cohomologies on symplectic manifolds that contain **unique harmonic representative within each class**:

PRIMITIVE COHOMOLOGIES [Tseng-Yau I & II, 12]

New finite dimensional cohomologies on symplectic manifolds that contain **unique harmonic representative within each class**:

- [Tseng-Yau I, 12] Idea: Bott-Chern and Aeppli in the symplectic setting.

$$H_{d+d^\wedge}^* = \frac{\text{Ker}(d + d^\wedge)}{\text{Im } dd^\wedge}, \quad H_{dd^\wedge}^* = \frac{\text{Ker } dd^\wedge}{\text{Im } d + \text{Im } d^\wedge}.$$

Primitive cohomologies: $PH_{d+d^\wedge}^q$ and $PH_{dd^\wedge}^q$, for $q \leq n$.

PRIMITIVE COHOMOLOGIES [Tseng-Yau I & II, 12]

New finite dimensional cohomologies on symplectic manifolds that contain **unique harmonic representative within each class**:

- [Tseng-Yau I, 12] Idea: Bott-Chern and Aeppli in the symplectic setting.

$$H_{d+d^\wedge}^* = \frac{\text{Ker}(d + d^\wedge)}{\text{Im } dd^\wedge}, \quad H_{dd^\wedge}^* = \frac{\text{Ker } dd^\wedge}{\text{Im } d + \text{Im } d^\wedge}.$$

Primitive cohomologies: $PH_{d+d^\wedge}^q$ and $PH_{dd^\wedge}^q$, for $q \leq n$.

- [Tseng-Yau II, 12] Idea: Dolbeault in symplectic case:

$$d = \partial_+ + \omega \wedge \partial_-$$

$$H_{\partial_+}^* \text{ and } H_{\partial_-}^*.$$

Primitive cohomologies: $PH_{\partial_+}^q$ and $PH_{\partial_-}^q$, for $q \leq n - 1$.

FILTERED COHOMOLOGIES [Tsai-Tseng-Yau III, 14]

$\{PH_{d+d^\wedge}^*, PH_{dd^\wedge}^*, PH_{\partial_+}^*, PH_{\partial_-}^*\}$ are primitive cohomologies.

FILTERED COHOMOLOGIES [Tsai-Tseng-Yau III, 14]

$\{PH_{d+d^\wedge}^*, PH_{dd^\wedge}^*, PH_{\partial_+}^*, PH_{\partial_-}^*\}$ are primitive cohomologies.

Idea: There exist non-primitive ones?

FILTERED COHOMOLOGIES [Tsai-Tseng-Yau III, 14]

$\{PH_{d+d^\wedge}^*, PH_{dd^\wedge}^*, PH_{\partial_+}^*, PH_{\partial_-}^*\}$ are primitive cohomologies.

Idea: There exist non-primitive ones?

Filtered cohomologies: $F^p H_+^q, F^p H_-^q$, for $q = 0, \dots, n + p$, and $p = 0, \dots, n$, that extend the primitive ones.

FILTERED COHOMOLOGIES [Tsai-Tseng-Yau III, 14]

$\{PH_{d+d^\wedge}^*, PH_{dd^\wedge}^*, PH_{\partial_+}^*, PH_{\partial_-}^*\}$ are primitive cohomologies.

Idea: There exist non-primitive ones?

Filtered cohomologies: $F^p H_+^q, F^p H_-^q$, for $q = 0, \dots, n + p$, and $p = 0, \dots, n$, that extend the primitive ones.

Relations between cohomologies for closed symplectic manifolds:

- $PH_{\partial_+}^q = PH_{\partial_-}^q, q \leq n - 1.$
- $PH_{dd^\wedge}^q = PH_{d+d^\wedge}^q, q \leq n.$
- $F^p H_+^{n+p} = PH_{dd^\wedge}^{n-p}.$
- $F^p H_-^{n+p} = PH_{d+d^\wedge}^{n-p}.$
- $F^p H_+^q = F^p H_-^q.$

COEFFECTIVE COHOMOLOGY I

Definition [Bouché 90]

$\alpha \in \Omega^*(M)$ is **coeffective** if $L_\omega(\alpha) := \alpha \wedge \omega = 0$. Notation: $\mathfrak{C}_{(1)}^q(M)$.

- $d(\mathfrak{C}_{(1)}^q(M)) \subset \mathfrak{C}_{(1)}^{q+1}(M)$.

Coeffective cohomology: $H_{(1)}^q(M) = \frac{\text{Ker} \{d : \mathfrak{C}_{(1)}^q(M) \rightarrow \mathfrak{C}_{(1)}^{q+1}(M)\}}{\text{Im} \{d : \mathfrak{C}_{(1)}^{q-1}(M) \rightarrow \mathfrak{C}_{(1)}^q(M)\}}$.

- $L_\omega : \Omega^q \rightarrow \Omega^{q+2}$ injective $\forall q \leq n-1 \implies H_{(1)}^q(M) = 0, \forall q \leq n-1$.
- $H_{(1)}^q(M) = H^q(M), q = 2n$.

COEFFECTIVE COHOMOLOGY II

If (M^{2n}, ω) is of finite type:

$$\underbrace{H_{(1)}^0(M), \dots, H_{(1)}^{n-1}(M)}_{=0}, H_{(1)}^n(M), \underbrace{H_{(1)}^{n+1}(M), \dots, H_{(1)}^{2n}(M)}_{\text{are finite dimensional}}.$$

Notation: $c_q^{(1)}(M) = \dim H_{(1)}^q(M)$, $q \geq n + 1$.

COEFFECTIVE COHOMOLOGY II

If (M^{2n}, ω) is of finite type:

$$\underbrace{H_{(1)}^0(M), \dots, H_{(1)}^{n-1}(M)}_{=0}, H_{(1)}^n(M), \underbrace{H_{(1)}^{n+1}(M), \dots, H_{(1)}^{2n}(M)}_{\text{are finite dimensional}}.$$

Notation: $c_q^{(1)}(M) = \dim H_{(1)}^q(M)$, $q \geq n + 1$.

Problem: $H_{(1)}^n(M)$ can be infinite dimensional

COEFFECTIVE COHOMOLOGY II

If (M^{2n}, ω) is of finite type:

$$\underbrace{H_{(1)}^0(M), \dots, H_{(1)}^{n-1}(M)}_{=0}, H_{(1)}^n(M), \underbrace{H_{(1)}^{n+1}(M), \dots, H_{(1)}^{2n}(M)}_{\text{are finite dimensional}}.$$

Notation: $c_q^{(1)}(M) = \dim H_{(1)}^q(M)$, $q \geq n + 1$.

Problem: $H_{(1)}^n(M)$ can be infinite dimensional

[Fernández-Ibáñez-de León 98] For $q \geq n + 1$:

$$b_q - b_{q+2} \leq c_q^{(1)} \leq b_q + b_{q+1}.$$

COEFFECTIVE COHOMOLOGY II

If (M^{2n}, ω) is of finite type:

$$\underbrace{H_{(1)}^0(M), \dots, H_{(1)}^{n-1}(M)}_{=0}, H_{(1)}^n(M), \underbrace{H_{(1)}^{n+1}(M), \dots, H_{(1)}^{2n}(M)}_{\text{are finite dimensional}}.$$

Notation: $c_q^{(1)}(M) = \dim H_{(1)}^q(M)$, $q \geq n + 1$.

Problem: $H_{(1)}^n(M)$ can be infinite dimensional

[Fernández-Ibáñez-de León 98] For $q \geq n + 1$:

$$b_q - b_{q+2} \leq c_q^{(1)} \leq b_q + b_{q+1}.$$

- HLC $\implies b_q - b_{q+2} = c_q^{(1)}$.

COEFFECTIVE COHOMOLOGY II

If (M^{2n}, ω) is of finite type:

$$\underbrace{H_{(1)}^0(M), \dots, H_{(1)}^{n-1}(M)}_{=0}, H_{(1)}^n(M), \underbrace{H_{(1)}^{n+1}(M), \dots, H_{(1)}^{2n}(M)}_{\text{are finite dimensional}}.$$

Notation: $c_q^{(1)}(M) = \dim H_{(1)}^q(M)$, $q \geq n + 1$.

Problem: $H_{(1)}^n(M)$ can be infinite dimensional

[Fernández-Ibáñez-de León 98] For $q \geq n + 1$:

$$b_q - b_{q+2} \leq c_q^{(1)} \leq b_q + b_{q+1}.$$

- HLC $\implies b_q - b_{q+2} = c_q^{(1)}$.
- ω exact $\implies b_q + b_{q+1} = c_q^{(1)}$.

k -COEFFEFFECTIVE COHOMOLOGIES I

$\alpha \in \Omega^*(M)$ is k -coeffective if $L_\omega^k(\alpha) := \alpha \wedge \omega^k = 0$. Notation: $e_{(k)}^q(M)$.

k -COEFFECTIVE COHOMOLOGIES I

$\alpha \in \Omega^*(M)$ is k -coeffective if $L_\omega^k(\alpha) := \alpha \wedge \omega^k = 0$. Notation: $\mathfrak{e}_{(k)}^q(M)$.

$$k\text{-Coeffective cohomology: } H_{(k)}^q(M) = \frac{\text{Ker } \{d : \mathfrak{e}_{(k)}^q(M) \rightarrow \mathfrak{e}_{(k)}^{q+1}(M)\}}{\text{Im } \{d : \mathfrak{e}_{(k)}^{q-1}(M) \rightarrow \mathfrak{e}_{(k)}^q(M)\}}.$$

k -COEFFECTIVE COHOMOLOGIES I

$\alpha \in \Omega^*(M)$ is **k -coeffective** if $L_\omega^k(\alpha) := \alpha \wedge \omega^k = 0$. Notation: $\mathfrak{e}_{(k)}^q(M)$.

k -Coeffective cohomology: $H_{(k)}^q(M) = \frac{\text{Ker} \{d : \mathfrak{e}_{(k)}^q(M) \rightarrow \mathfrak{e}_{(k)}^{q+1}(M)\}}{\text{Im} \{d : \mathfrak{e}_{(k)}^{q-1}(M) \rightarrow \mathfrak{e}_{(k)}^q(M)\}}$.

If (M^{2n}, ω) is of finite type:

$$\underbrace{H_{(k)}^0, \dots, H_{(k)}^{n-k}}_{=0}, H_{(k)}^{n-k+1}, H_{(k)}^{n-k+2}, \dots, H_{(k)}^{2n-2k+1}, \underbrace{H_{(k)}^{2n-2k+2}, \dots, H_{(k)}^{2n}}_{=H^q}.$$

are finite dimensional

k -COEFFECTIVE COHOMOLOGIES I

$\alpha \in \Omega^*(M)$ is **k -coeffective** if $L_\omega^k(\alpha) := \alpha \wedge \omega^k = 0$. Notation: $\mathfrak{e}_{(k)}^q(M)$.

k -Coeffective cohomology: $H_{(k)}^q(M) = \frac{\text{Ker} \{d : \mathfrak{e}_{(k)}^q(M) \rightarrow \mathfrak{e}_{(k)}^{q+1}(M)\}}{\text{Im} \{d : \mathfrak{e}_{(k)}^{q-1}(M) \rightarrow \mathfrak{e}_{(k)}^q(M)\}}$.

If (M^{2n}, ω) is of finite type:

$$\underbrace{H_{(k)}^0, \dots, H_{(k)}^{n-k}}_{=0}, H_{(k)}^{n-k+1}, H_{(k)}^{n-k+2}, \dots, H_{(k)}^{2n-2k+1}, \underbrace{H_{(k)}^{2n-2k+2}, \dots, H_{(k)}^{2n}}_{=H^q}.$$

are finite dimensional

Problem: $H_{(k)}^{n-k+1}(M)$ can be infinite dimensional

k -COEFFEFFECTIVE COHOMOLOGIES II

Notation: $c_q^{(k)}(M) = \dim H_{(k)}^q(M)$, $q \geq n - k + 2$.

k -COEFFECTIVE COHOMOLOGIES II

Notation: $c_q^{(k)}(M) = \dim H_{(k)}^q(M)$, $q \geq n - k + 2$.

$$b_q - b_{q+2k} \leq c_q^{(k)} \leq b_q + b_{q+2k-1}. \quad (1)$$

- HLC $\implies b_q - b_{q+2k} = c_q^{(k)}$.
- ω exact $\implies b_q + b_{q+2k-1} = c_q^{(k)}$.

k -COEFFECTIVE COHOMOLOGIES II

Notation: $c_q^{(k)}(M) = \dim H_{(k)}^q(M)$, $q \geq n - k + 2$.

$$b_q - b_{q+2k} \leq c_q^{(k)} \leq b_q + b_{q+2k-1}. \quad (1)$$

- HLC $\implies b_q - b_{q+2k} = c_q^{(k)}$.
- ω exact $\implies b_q + b_{q+2k-1} = c_q^{(k)}$.

Objective

Define a new group for degree $n - k + 1$ such that:

- It is finite dimensional.
- Its dimension satisfies similar inequalities to (1)

GENERALIZED COEFFECTIVE COHOMOLOGIES

If (M^{2n}, ω) is of finite type, for degree $n - k + 1$ we define a new finite-dimensional space (using a long exact sequence in cohomology)

Definition

$$\hat{H}^{n-k+1}(M) = \frac{H_{(k)}^{n-k+1}(M)}{\frac{H^{n+k}(L_{\omega}^k(\Omega^k(M)))}{H^{n-k}(M)}}. \quad \dim \hat{H}^{n-k+1}(M) = \hat{c}_{n-k+1}.$$

- $b_{n-k+1} - b_{n+k+1} \leq \hat{c}_{n-k+1} \leq b_{n-k+1}$ (HLC and exact \Rightarrow)

GENERALIZED COEFFECTIVE COHOMOLOGIES

If (M^{2n}, ω) is of finite type, for degree $n - k + 1$ we define a new finite-dimensional space (using a long exact sequence in cohomology)

Definition

$$\hat{H}^{n-k+1}(M) = \frac{H_{(k)}^{n-k+1}(M)}{\frac{H^{n+k}(L_{\omega}^k(\Omega^k(M)))}{H^{n-k}(M)}}. \quad \dim \hat{H}^{n-k+1}(M) = \hat{c}_{n-k+1}.$$

- $b_{n-k+1} - b_{n+k+1} \leq \hat{c}_{n-k+1} \leq b_{n-k+1}$ (HLC and exact \Rightarrow)

Generalized coeffective cohomology

$$\hat{H}^{n-k+1}, H_{(k)}^{n-k+2}, \dots, H_{(k)}^{2n}, \quad 1 \leq k \leq n.$$

GENERALIZED COEFFECTIVE COHOMOLOGIES

If (M^{2n}, ω) is of finite type, for degree $n - k + 1$ we define a new finite-dimensional space (using a long exact sequence in cohomology)

Definition

$$\hat{H}^{n-k+1}(M) = \frac{H_{(k)}^{n-k+1}(M)}{\frac{H^{n+k}(L_{\omega}^k(\Omega^k(M)))}{H^{n-k}(M)}}. \quad \dim \hat{H}^{n-k+1}(M) = \hat{c}_{n-k+1}.$$

- $b_{n-k+1} - b_{n+k+1} \leq \hat{c}_{n-k+1} \leq b_{n-k+1}$ (HLC and exact \Rightarrow)

Generalized coeffective cohomology

$$\hat{H}^{n-k+1}, H_{(k)}^{n-k+2}, \dots, H_{(k)}^{2n}, \quad 1 \leq k \leq n.$$

- $\chi^{(k)}(M) = (-1)^{n-k+1} \hat{c}_{n-k+1} + \sum_{i=n-k+2}^{2n} (-1)^i c_i^{(k)}$

GENERALIZED COEFFECTIVE COHOMOLOGIES

If (M^{2n}, ω) is of finite type, for degree $n - k + 1$ we define a new finite-dimensional space (using a long exact sequence in cohomology)

Definition

$$\hat{H}^{n-k+1}(M) = \frac{H_{(k)}^{n-k+1}(M)}{\frac{H^{n+k}(L_{\omega}^k(\Omega^k(M)))}{H^{n-k}(M)}}. \quad \dim \hat{H}^{n-k+1}(M) = \hat{c}_{n-k+1}.$$

- $b_{n-k+1} - b_{n+k+1} \leq \hat{c}_{n-k+1} \leq b_{n-k+1}$ (HLC and exact \Rightarrow)

Generalized coeffective cohomology

$$\hat{H}^{n-k+1}, H_{(k)}^{n-k+2}, \dots, H_{(k)}^{2n}, \quad 1 \leq k \leq n.$$

- $\chi^{(k)}(M) = (-1)^{n-k+1} \hat{c}_{n-k+1} + \sum_{i=n-k+2}^{2n} (-1)^i c_i^{(k)} = \sum_{i=n-k+1}^{n+k} (-1)^i b_i$

GENERALIZED COEFFECTIVE COHOMOLOGIES

If (M^{2n}, ω) is of finite type, for degree $n - k + 1$ we define a new finite-dimensional space (using a long exact sequence in cohomology)

Definition

$$\hat{H}^{n-k+1}(M) = \frac{H_{(k)}^{n-k+1}(M)}{\frac{H^{n+k}(L_{\omega}^k(\Omega^k(M)))}{H^{n-k}(M)}}. \quad \dim \hat{H}^{n-k+1}(M) = \hat{c}_{n-k+1}.$$

- $b_{n-k+1} - b_{n+k+1} \leq \hat{c}_{n-k+1} \leq b_{n-k+1}$ (HLC and exact \Rightarrow)

Generalized coeffective cohomology

$$\hat{H}^{n-k+1}, H_{(k)}^{n-k+2}, \dots, H_{(k)}^{2n}, \quad 1 \leq k \leq n.$$

- $\chi^{(k)}(M) = (-1)^{n-k+1} \hat{c}_{n-k+1} + \sum_{i=n-k+2}^{2n} (-1)^i c_i^{(k)} = \sum_{i=n-k+1}^{n+k} (-1)^i b_i$

topological invariant.

RELATIONS BETWEEN COEFFECTIVE AND HARMONIC COHOMOLOGIES

Theorem

Let (M^{2n}, ω) be a symplectic manifold of finite type. The following relation holds for every $k = 1, \dots, n$:

$$h_{n-k+1} - h_{n+k+1} = \hat{c}_{n-k+1}.$$

RELATIONS BETWEEN COEFFECTIVE AND HARMONIC COHOMOLOGIES

Theorem

Let (M^{2n}, ω) be a symplectic manifold of finite type. The following relation holds for every $k = 1, \dots, n$:

$$h_{n-k+1} - h_{n+k+1} = \hat{c}_{n-k+1}.$$

$$h_0, h_1, h_2, \dots, \overbrace{h_n, h_{n+1}, h_{n+2}, \dots, h_{2n}}^{\hat{c}_1}, \dots, \overbrace{h_0, h_1, h_2, \dots, h_n, h_{n+1}, h_{n+2}, \dots, h_{2n}}^{\hat{c}_{n-1}}$$

No relation for h_{n+1} .

EXTENSIONS OF THE GENERALIZED COEFFECTIVE COMPLEXES I

The coeffective complexes are not elliptic in degree $n - k + 1$ (that is the reason for which $H_{(k)}^{n-k+1}$ can be infinite dimensional).

EXTENSIONS OF THE GENERALIZED COEFFECTIVE COMPLEXES I

The coeffective complexes are not elliptic in degree $n - k + 1$ (that is the reason for which $H_{(k)}^{n-k+1}$ can be infinite dimensional).

Idea: Construct an **elliptic** extension of the coeffective complexes.
([Eastwood 12] for $k = 1$.)

EXTENSIONS OF THE GENERALIZED COEFFECTIVE COMPLEXES I

The coeffective complexes are not elliptic in degree $n - k + 1$ (that is the reason for which $H_{(k)}^{n-k+1}$ can be infinite dimensional).

Idea: Construct an **elliptic** extension of the coeffective complexes.
 ([Eastwood 12] for $k = 1$.)

$$\begin{array}{cccccccccccc}
 0 & \longrightarrow & \Omega^0 & \xrightarrow{d} & \dots & \xrightarrow{d} & \Omega^{2k-1} & \xrightarrow{\check{d}} & \check{\Omega}_{(k)}^{2k} & \xrightarrow{\check{d}} & \dots & \xrightarrow{\check{d}} & \check{\Omega}_{(k)}^{n+k-2} & \xrightarrow{\check{d}} & \check{\Omega}_{(k)}^{n+k-1} \\
 & & & & & & & & & & & & & & & \downarrow D \\
 0 & \longleftarrow & \Omega^{2n} & \xleftarrow{d} & \dots & \xleftarrow{d} & \Omega^{2n-2k+1} & \xleftarrow{d} & \mathfrak{C}_{(k)}^{2n-2k} & \xleftarrow{d} & \dots & \xleftarrow{d} & \mathfrak{C}_{(k)}^{n-k+2} & \xleftarrow{d} & \mathfrak{C}_{(k)}^{n-k+1}
 \end{array}$$

where $\check{\Omega}_{(k)}^q(M) = \frac{\Omega^q(M)}{L_{\omega}^k(\Omega^{q-2k}(M))}$, \check{d} is induced by d , and D is second order operator.

EXTENSIONS OF THE GENERALIZED COEFFECTIVE COMPLEXES II

Cohomology groups: $\check{H}_{(k)}^q(M)$, for $q = 0, \dots, 2n + 2k - 1$.

EXTENSIONS OF THE GENERALIZED COEFFECTIVE COMPLEXES II

Cohomology groups: $\check{H}_{(k)}^q(M)$, for $q = 0, \dots, 2n + 2k - 1$.

$$\underbrace{\check{H}_{(k)}^0, \dots, \check{H}_{(k)}^{2k-2}}_{= H^q}, \check{H}_{(k)}^{2k-1}, \dots, \check{H}_{(k)}^{n+k}, \underbrace{\check{H}_{(k)}^{n+k+1}, \dots, \check{H}_{(k)}^{2n+2k-1}}_{= H_{(k)}^{q-2k+1}}.$$

EXTENSIONS OF THE GENERALIZED COEFFECTIVE COMPLEXES II

Cohomology groups: $\check{H}_{(k)}^q(M)$, for $q = 0, \dots, 2n + 2k - 1$.

$$\underbrace{\check{H}_{(k)}^0, \dots, \check{H}_{(k)}^{2k-2}, \check{H}_{(k)}^{2k-1}, \dots, \check{H}_{(k)}^{n+k}}_{= H^q}, \quad \underbrace{\check{H}_{(k)}^{n+k+1}, \dots, \check{H}_{(k)}^{2n+2k-1}}_{= H_{(k)}^{q-2k+1}}.$$

- If (M^{2n}, ω) is of finite type, $\check{c}_q^{(k)}(M) = \dim \check{H}_{(k)}^q(M)$ finite $\forall q$.

EXTENSIONS OF THE GENERALIZED COEFFECTIVE COMPLEXES II

Cohomology groups: $\check{H}_{(k)}^q(M)$, for $q = 0, \dots, 2n + 2k - 1$.

$$\underbrace{\check{H}_{(k)}^0, \dots, \check{H}_{(k)}^{2k-2}, \check{H}_{(k)}^{2k-1}, \dots, \check{H}_{(k)}^{n+k}}_{= H^q}, \quad \underbrace{\check{H}_{(k)}^{n+k+1}, \dots, \check{H}_{(k)}^{2n+2k-1}}_{= H_{(k)}^{q-2k+1}}.$$

- If (M^{2n}, ω) is of finite type, $\check{c}_q^{(k)}(M) = \dim \check{H}_{(k)}^q(M)$ finite $\forall q$.

$$b_{q-2k+1} - b_{q+1} \leq \check{c}_q^{(k)} \leq b_{q-2k+1} + b_q.$$

EXTENSIONS OF THE GENERALIZED COEFFECTIVE COMPLEXES II

Cohomology groups: $\check{H}_{(k)}^q(M)$, for $q = 0, \dots, 2n + 2k - 1$.

$$\underbrace{\check{H}_{(k)}^0, \dots, \check{H}_{(k)}^{2k-2}, \check{H}_{(k)}^{2k-1}, \dots, \check{H}_{(k)}^{n+k}}_{= H^q}, \quad \underbrace{\check{H}_{(k)}^{n+k+1}, \dots, \check{H}_{(k)}^{2n+2k-1}}_{= H_{(k)}^{q-2k+1}}.$$

- If (M^{2n}, ω) is of finite type, $\check{c}_q^{(k)}(M) = \dim \check{H}_{(k)}^q(M)$ finite $\forall q$.

$$b_{q-2k+1} - b_{q+1} \leq \check{c}_q^{(k)} \leq b_{q-2k+1} + b_q.$$

- ▶ HLC $\implies b_{q-2k+1} - b_{q+1} = \check{c}_q^{(k)}$.
- ▶ ω exact $\implies b_{q-2k+1} + b_q = \check{c}_q^{(k)}$.

RELATIONS BETWEEN COEFFECTIVE AND PRIMITIVE COHOMOLOGIES

These last cohomology groups allow us to recover all the **primitive** cohomology groups [Tseng-Yau 12] via the filtered cohomology groups [Tsai-Tseng-Yau 14]:

RELATIONS BETWEEN COEFFECTIVE AND PRIMITIVE COHOMOLOGIES

These last cohomology groups allow us to recover all the **primitive** cohomology groups [Tseng-Yau 12] via the filtered cohomology groups [Tsai-Tseng-Yau 14]:

$$PH_{\partial_+}^q(M) \cong F^0 H_+^q(M) \cong \check{H}_{(1)}^q(M), \quad 0 \leq q \leq n-1;$$

$$PH_{\partial_-}^q(M) \cong F^0 H_-^q(M) \cong \check{H}_{(1)}^{2n-q+1}(M) \cong H_{(1)}^{2n-q}(M), \quad 0 \leq q \leq n-1;$$

$$PH_{dd^\Lambda}^{n-k+1}(M) \cong F^{k-1} H_+^{n+k-1}(M) \cong \check{H}_{(k)}^{n+k-1}(M), \quad 1 \leq k \leq n;$$

$$PH_{d+d^\Lambda}^{n-k+1}(M) \cong F^{k-1} H_-^{n+k-1}(M) \cong \check{H}_{(k)}^{n+k}(M), \quad 1 \leq k \leq n.$$

RELATIONS BETWEEN COEFFECTIVE AND PRIMITIVE COHOMOLOGIES

These last cohomology groups allow us to recover all the **primitive** cohomology groups [Tseng-Yau 12] via the filtered cohomology groups [Tsai-Tseng-Yau 14]:

$$PH_{\partial_+}^q(M) \cong F^0 H_+^q(M) \cong \check{H}_{(1)}^q(M), \quad 0 \leq q \leq n-1;$$

$$PH_{\partial_-}^q(M) \cong F^0 H_-^q(M) \cong \check{H}_{(1)}^{2n-q+1}(M) \cong H_{(1)}^{2n-q}(M), \quad 0 \leq q \leq n-1;$$

$$PH_{dd^{\wedge}}^{n-k+1}(M) \cong F^{k-1} H_+^{n+k-1}(M) \cong \check{H}_{(k)}^{n+k-1}(M), \quad 1 \leq k \leq n;$$

$$PH_{d+d^{\wedge}}^{n-k+1}(M) \cong F^{k-1} H_-^{n+k-1}(M) \cong \check{H}_{(k)}^{n+k}(M), \quad 1 \leq k \leq n.$$

In fact, the following isomorphisms hold:

$$F^{k-1} H_-^{n+k-s-1}(M) \cong H_{(k)}^{n-k+s+1}(M) \cong \check{H}_{(k)}^{n+k+s}(M), \quad 1 \leq s \leq n+k-1.$$

RELATIONS BETWEEN COEFFECTIVE AND PRIMITIVE COHOMOLOGIES

These last cohomology groups allow us to recover all the **primitive** cohomology groups [Tseng-Yau 12] via the filtered cohomology groups [Tsai-Tseng-Yau 14]:

$$PH_{\partial_+}^q(M) \cong F^0 H_+^q(M) \cong \check{H}_{(1)}^q(M), \quad 0 \leq q \leq n-1;$$

$$PH_{\partial_-}^q(M) \cong F^0 H_-^q(M) \cong \check{H}_{(1)}^{2n-q+1}(M) \cong H_{(1)}^{2n-q}(M), \quad 0 \leq q \leq n-1;$$

$$PH_{dd^{\wedge}}^{n-k+1}(M) \cong F^{k-1} H_+^{n+k-1}(M) \cong \check{H}_{(k)}^{n+k-1}(M), \quad 1 \leq k \leq n;$$

$$PH_{d+d^{\wedge}}^{n-k+1}(M) \cong F^{k-1} H_-^{n+k-1}(M) \cong \check{H}_{(k)}^{n+k}(M), \quad 1 \leq k \leq n.$$

In fact, the following isomorphisms hold:

$$F^{k-1} H_-^{n+k-s-1}(M) \cong H_{(k)}^{n-k+s+1}(M) \cong \check{H}_{(k)}^{n+k+s}(M), \quad 1 \leq s \leq n+k-1.$$

$$\check{H}_{(k)}^{n+k+s}(M) \cong \check{H}_{(k)}^{n+k-s-1}(M) \text{ for } s = 0, \dots, n+k-1.$$

MORE RELATIONS BETWEEN COHOMOLOGIES

Let (M^{2n}, ω) be symplectic of finite type.

$$0 \leq \check{c}_{n+k}^{(k)} - \hat{c}_{n-k+1} \leq b_{n+k} - h_{n+k} \quad (2)$$

MORE RELATIONS BETWEEN COHOMOLOGIES

Let (M^{2n}, ω) be symplectic of finite type.

$$0 \leq \check{c}_{n+k}^{(k)} - \hat{c}_{n-k+1} \leq b_{n+k} - h_{n+k} \quad (2)$$

$$\check{c}_{n+k}^{(k)} - \hat{c}_{n-k+1} = b_{n+k} - h_{n+k} \iff L^k(H_{\text{hr}}^{n-k}(M)) = L^k(H^{n-k}(M)).$$

MORE RELATIONS BETWEEN COHOMOLOGIES

Let (M^{2n}, ω) be symplectic of finite type.

$$0 \leq \check{c}_{n+k}^{(k)} - \hat{c}_{n-k+1} \leq b_{n+k} - h_{n+k} \quad (2)$$

$$\check{c}_{n+k}^{(k)} - \hat{c}_{n-k+1} = b_{n+k} - h_{n+k} \iff L^k(H_{\text{hr}}^{n-k}(M)) = L^k(H^{n-k}(M)).$$

Particular cases for equality: $k = n, n - 1, n - 2 \implies$

- $\check{c}_{2n}^{(n)} = b_1 + b_{2n} - h_{2n}.$
- $\check{c}_{2n-1}^{(n-1)} = b_2 + b_{2n-1} - h_{2n-1} - h_{2n}.$
- $\check{c}_{2n-2}^{(n-2)} = b_{2n-2} + h_3 - h_{2n-2} - h_{2n-1}.$

MORE RELATIONS BETWEEN COHOMOLOGIES

Let (M^{2n}, ω) be symplectic of finite type.

$$0 \leq \check{c}_{n+k}^{(k)} - \hat{c}_{n-k+1} \leq b_{n+k} - h_{n+k} \quad (2)$$

$$\check{c}_{n+k}^{(k)} - \hat{c}_{n-k+1} = b_{n+k} - h_{n+k} \iff L^k(H_{\text{hr}}^{n-k}(M)) = L^k(H^{n-k}(M)).$$

Particular cases for equality: $k = n, n - 1, n - 2 \implies$

- $\check{c}_{2n}^{(n)} = b_1 + b_{2n} - h_{2n}.$
- $\check{c}_{2n-1}^{(n-1)} = b_2 + b_{2n-1} - h_{2n-1} - h_{2n}.$
- $\check{c}_{2n-2}^{(n-2)} = b_{2n-2} + h_3 - h_{2n-2} - h_{2n-1}.$

Theorem

(M^{2n}, ω) satisfies **HLC** $\iff \check{c}_{n+k}^{(k)} = \hat{c}_{n-k+1}$ for $k = 1, \dots, n.$

RELATIONS FOR CLOSED SYMPLECTIC MANIFOLDS

k -generalized coeffective cohomologies

- $k = 1, \dots, n - 1 : c_q^{(k)} = b_q, \quad \text{for } q = 2n - 2k + 1, \dots, 2n.$
- $k = n : \begin{cases} c_q^{(n)} = b_q, & q = 2, \dots, 2n, \\ \hat{c}_1 = b_1. \end{cases}$
- $\chi^{(k)}.$

RELATIONS FOR CLOSED SYMPLECTIC MANIFOLDS

k -generalized coeffective cohomologies

- $k = 1, \dots, n-1$: $c_q^{(k)} = b_q$, for $q = 2n - 2k + 1, \dots, 2n$.
- $k = n$: $\begin{cases} c_q^{(n)} = b_q, & q = 2, \dots, 2n, \\ \hat{c}_1 = b_1. \end{cases}$
- $\chi^{(k)}$.

filtered cohomologies

$$\check{c}_q^{(k)} = \begin{cases} \check{c}_{2n+2k-q-1}^{(k)}, & \text{if } q = 0, \dots, n+k-1, \\ c_{q-2k+1}^{(k)}, & \text{if } q = n+k+1, \dots, 2n+2k-1. \end{cases}$$

Explicit descriptions of $\check{c}_{2n}^{(n)}$, $\check{c}_{2n-1}^{(n-1)}$, $\check{c}_{2n-2}^{(n-2)}$.

RELATIONS FOR CLOSED SYMPLECTIC MANIFOLDS

k -generalized coeffective cohomologies

- $k = 1, \dots, n-1$: $c_q^{(k)} = b_q$, for $q = 2n - 2k + 1, \dots, 2n$.
- $k = n$: $\begin{cases} c_q^{(n)} = b_q, & q = 2, \dots, 2n, \\ \hat{c}_1 = b_1. \end{cases}$ topological for $k = n, n-1$.
- $\chi^{(k)}$.

filtered cohomologies

$$\check{c}_q^{(k)} = \begin{cases} \check{c}_{2n+2k-q-1}^{(k)}, & \text{if } q = 0, \dots, n+k-1, \\ c_{q-2k+1}^{(k)}, & \text{if } q = n+k+1, \dots, 2n+2k-1. \end{cases} \quad \text{top for } k = n.$$

Explicit descriptions of $\check{c}_{2n}^{(n)}$, $\check{c}_{2n-1}^{(n-1)}$, $\check{c}_{2n-2}^{(n-2)}$.

RELATIONS FOR CLOSED SYMPLECTIC MANIFOLDS

k -generalized coeffective cohomologies

- $k = 1, \dots, n-1 : c_q^{(k)} = b_q$, for $q = 2n - 2k + 1, \dots, 2n$.
- $k = n : \begin{cases} c_q^{(n)} = b_q, & q = 2, \dots, 2n, \\ \hat{c}_1 = b_1. \end{cases}$ topological for $k = n, n-1$.
- $\chi^{(k)}$.

filtered cohomologies

$$\check{c}_q^{(k)} = \begin{cases} \check{c}_{2n+2k-q-1}^{(k)}, & \text{if } q = 0, \dots, n+k-1, \\ c_{q-2k+1}^{(k)}, & \text{if } q = n+k+1, \dots, 2n+2k-1. \end{cases}$$

top for $k = n$.

Explicit descriptions of $\check{c}_{2n}^{(n)}$, $\check{c}_{2n-1}^{(n-1)}$, $\check{c}_{2n-2}^{(n-2)}$.

harmonic cohomology

- $h_i = b_i$, $i = 0, 1, 2, 2n$, h_{2n-1} even.
- $h_{n-k+1} - h_{n+k+1} = \hat{c}_{n-k+1}$. $h_{n-1} - (\check{c}_{n+1}^{(1)} - \hat{c}_n) \leq h_{n+1} \leq h_{n-1}$.

FLEXIBILITY

An interesting question in the study of the symplectic harmonicity is the **flexibility**. It was introduced and studied in [IRTU 01] and [Yan 96] motivated by a question posed by Khesin and McDuff:

FLEXIBILITY

An interesting question in the study of the symplectic harmonicity is the **flexibility**. It was introduced and studied in [IRTU 01] and [Yan 96] motivated by a question posed by Khesin and McDuff:

Question: On which compact manifolds M does there exist a family ω_t of symplectic forms such that the dimension of $H_{\text{hr}}^q(M, \omega_t)$ varies?

FLEXIBILITY

An interesting question in the study of the symplectic harmonicity is the **flexibility**. It was introduced and studied in [IRTU 01] and [Yan 96] motivated by a question posed by Khesin and McDuff:

Question: On which compact manifolds M does there exist a family ω_t of symplectic forms such that the dimension of $H_{\text{hr}}^q(M, \omega_t)$ varies?

Definition

A manifold M^{2n} is **h-flexible** if M possesses a continuous symplectic family ω_t , $t \in [a, b]$ s.t. $h_q(M, \omega_a) \neq h_q(M, \omega_b)$ for some q .

FLEXIBILITY

An interesting question in the study of the symplectic harmonicity is the **flexibility**. It was introduced and studied in [IRTU 01] and [Yan 96] motivated by a question posed by Khesin and McDuff:

Question: On which compact manifolds M does there exist a family ω_t of symplectic forms such that the dimension of $H_{\text{hr}}^q(M, \omega_t)$ varies?

Definition

A manifold M^{2n} is **h-flexible** if M possesses a continuous symplectic family ω_t , $t \in [a, b]$ s.t. $h_q(M, \omega_a) \neq h_q(M, \omega_b)$ for some q .

- [Yan 96] studied the case of closed 4-manifolds:
 - ▶ 4-dimensional nilmanifolds (compact quotients of nilpotent Lie groups) are not **h-flexible**.
 - ▶ There exist 4-dimensional **h-flexible** manifolds.

FLEXIBILITY

An interesting question in the study of the symplectic harmonicity is the **flexibility**. It was introduced and studied in [IRTU 01] and [Yan 96] motivated by a question posed by Khesin and McDuff:

Question: On which compact manifolds M does there exist a family ω_t of symplectic forms such that the dimension of $H_{\text{hr}}^q(M, \omega_t)$ varies?

Definition

A manifold M^{2n} is **h-flexible** if M possesses a continuous symplectic family ω_t , $t \in [a, b]$ s.t. $h_q(M, \omega_a) \neq h_q(M, \omega_b)$ for some q .

- [Yan 96] studied the case of closed 4-manifolds:
 - ▶ 4-dimensional nilmanifolds (compact quotients of nilpotent Lie groups) are not **h-flexible**.
 - ▶ There exist 4-dimensional **h-flexible** manifolds.
- [IRTU 01]: Some 6-dimensional nilmanifolds are **h-flexible**.

COHOMOLOGICAL FLEXIBILITIES

We can define other notions of flexibility:

COHOMOLOGICAL FLEXIBILITIES

We can define other notions of flexibility:

Definition

A closed smooth manifold M^{2n} is said to be

- ① **c-flexible** if M possesses a continuous symplectic family ω_t , $t \in [a, b]$ such that, for some $1 \leq k \leq n$

$$\begin{cases} \hat{c}_{n-k+1}(M, \omega_a) \neq \hat{c}_{n-k+1}(M, \omega_b), & \text{or} \\ c_q^{(k)}(M, \omega_a) \neq c_q^{(k)}(M, \omega_b), & \text{for some } n - k + 2 \leq q \leq 2n. \end{cases}$$

- ② **f-flexible** if M possesses a continuous symplectic family ω_t , $t \in [a, b]$ such that $\check{c}_q^{(k)}(M, \omega_a) \neq \check{c}_q^{(k)}(M, \omega_b)$ for some $1 \leq k \leq n$ and $0 \leq q \leq 2n + 2k - 1$.

COHOMOLOGICAL FLEXIBILITIES

We can define other notions of flexibility:

Definition

A closed smooth manifold M^{2n} is said to be

- ① **c-flexible** if M possesses a continuous symplectic family ω_t , $t \in [a, b]$ such that, for some $1 \leq k \leq n$

$$\begin{cases} \hat{c}_{n-k+1}(M, \omega_a) \neq \hat{c}_{n-k+1}(M, \omega_b), & \text{or} \\ c_q^{(k)}(M, \omega_a) \neq c_q^{(k)}(M, \omega_b), & \text{for some } n - k + 2 \leq q \leq 2n. \end{cases}$$

- ② **f-flexible** if M possesses a continuous symplectic family ω_t , $t \in [a, b]$ such that $\check{c}_q^{(k)}(M, \omega_a) \neq \check{c}_q^{(k)}(M, \omega_b)$ for some $1 \leq k \leq n$ and $0 \leq q \leq 2n + 2k - 1$.

These new notions of flexibility give us a simpler way to provide examples of **h-flexible** manifolds.

CLOSED 4-DIMENSIONAL SYMPLECTIC MANIFOLDS

Let (M^4, ω) be a closed symplectic manifold.

Generalized coeffective cohomologies are topological.

CLOSED 4-DIMENSIONAL SYMPLECTIC MANIFOLDS

Let (M^4, ω) be a closed symplectic manifold.

Generalized coeffective cohomologies are topological.

Filtered cohomologies ($k = 2$ is topological)

$$k = 1: \check{c}_0^{(1)}, \check{c}_1^{(1)}, \check{c}_2^{(1)} \mid \check{c}_3^{(1)}, \underbrace{\check{c}_4^{(1)}}_{=c_3^{(1)}}, \underbrace{\check{c}_5^{(1)}}_{=c_4^{(1)}}, \text{ where } \boxed{\check{c}_3^{(1)} = b_1 + b_2 - h_3 - 1}$$

CLOSED 4-DIMENSIONAL SYMPLECTIC MANIFOLDS

Let (M^4, ω) be a closed symplectic manifold.

Generalized coeffective cohomologies are topological.

Filtered cohomologies ($k = 2$ is topological)

$$k = 1: \check{c}_0^{(1)}, \check{c}_1^{(1)}, \check{c}_2^{(1)} \mid \check{c}_3^{(1)}, \underbrace{\check{c}_4^{(1)}}_{=c_3^{(1)}}, \underbrace{\check{c}_5^{(1)}}_{=c_4^{(1)}}, \text{ where } \boxed{\check{c}_3^{(1)} = b_1 + b_2 - h_3 - 1}$$

$$\text{Harmonic cohomology: } \underbrace{h_0}_{=1}, \underbrace{h_1}_{=b_1}, \underbrace{h_2}_{=b_2}, \underbrace{h_3}_{\text{even}}, \underbrace{h_4}_{=1}.$$

CLOSED 4-DIMENSIONAL SYMPLECTIC MANIFOLDS

Let (M^4, ω) be a closed symplectic manifold.

Generalized coeffective cohomologies are topological.

Filtered cohomologies ($k = 2$ is topological)

$$k = 1: \check{c}_0^{(1)}, \check{c}_1^{(1)}, \check{c}_2^{(1)} \mid \check{c}_3^{(1)}, \underbrace{\check{c}_4^{(1)}}_{=c_3^{(1)}}, \underbrace{\check{c}_5^{(1)}}_{=c_4^{(1)}}, \text{ where } \check{c}_3^{(1)} = b_1 + b_2 - h_3 - 1$$

$$\text{Harmonic cohomology: } \underbrace{h_0}_{=1}, \underbrace{h_1}_{=b_1}, \underbrace{h_2}_{=b_2}, \underbrace{h_3}_{\text{even}}, \underbrace{h_4}_{=1}.$$

- M is never **c**-flexible.
- M is **f**-flexible $\iff M$ is **h**-flexible. In particular, $\exists M^4$ **f**-flexible.
- If $b_1(M) \leq 1$, then M is not **f**-flexible.
- If M is completely solvable solvmanifold, it is not **c**-flexible, **f**-flexible or **h**-flexible.

CLOSED 6-DIMENSIONAL SYMPLECTIC MANIFOLDS

Generalized coeffective cohomologies

$k = 1$:

$$\hat{c}_3, c_4^{(1)}, \underbrace{c_5^{(1)}}_{=b_5}, \underbrace{c_6^{(1)}}_{=b_6},$$

where

$$\hat{c}_3 = c_4^{(1)} + 1 - b_1 - b_2 + b_3$$

$k = 2$ and $k = 3$ are topological.

CLOSED 6-DIMENSIONAL SYMPLECTIC MANIFOLDS

Generalized coeffective cohomologies

$k = 1$:

$$\hat{c}_3, c_4^{(1)}, \underbrace{c_5^{(1)}}_{=b_5}, \underbrace{c_6^{(1)}}_{=b_6},$$

where

$$\hat{c}_3 = c_4^{(1)} + 1 - b_1 - b_2 + b_3$$

$k = 2$ and $k = 3$ are topological.

Harmonic cohomology

$$\underbrace{h_0}_{=1}, \underbrace{h_1}_{=b_1}, \underbrace{h_2}_{=b_2}, h_3, h_4, \underbrace{h_5}_{\text{even}}, \underbrace{h_6}_{=1},$$

where

$$\hat{c}_3 = h_3 - h_5$$

CLOSED 6-DIMENSIONAL SYMPLECTIC MANIFOLDS

Filtered cohomologies ($k = 3$ is topological)

$k = 1$:

$$\check{c}_0^{(1)}, \check{c}_1^{(1)}, \check{c}_2^{(1)}, \check{c}_3^{(1)} \mid \check{c}_4^{(1)}, \underbrace{\check{c}_5^{(1)}}_{=c_4^{(1)}}, \underbrace{\check{c}_6^{(1)}}_{=b_5}, \underbrace{\check{c}_7^{(1)}}_{=b_6},$$

where $\boxed{\check{c}_4^{(1)} = \hat{c}_3 - h_4 + b_2, \quad \check{c}_5^{(1)} = c_4^{(1)}}$

$k = 2$:

$$\check{c}_0^{(2)}, \check{c}_1^{(2)}, \check{c}_2^{(2)}, \check{c}_3^{(2)}, \check{c}_4^{(2)} \mid \check{c}_5^{(2)}, \underbrace{\check{c}_6^{(2)}}_{=b_3}, \underbrace{\check{c}_7^{(2)}}_{=b_4}, \underbrace{\check{c}_8^{(2)}}_{=b_5}, \underbrace{\check{c}_9^{(2)}}_{=b_6},$$

where $\boxed{\check{c}_5^{(2)} = b_1 + b_2 - h_5 - 1}$

All the dimensions are determined by **Betti numbers** and \hat{c}_3 , h_4 and h_5 .

FLEXIBILITY IN DIMENSION 6

Relations:

$$\begin{aligned}\hat{c}_3 &= c_4^{(1)} + 1 - b_1 - b_2 + b_3, \\ \hat{c}_3 &= h_3 - h_5, \\ \check{c}_4^{(1)} &= \hat{c}_3 - h_4 + b_2, \\ \check{c}_5^{(1)} &= c_4^{(1)}, \\ \check{c}_5^{(2)} &= b_1 + b_2 - h_5 - 1.\end{aligned}$$

Results:

- 1 M is **c**-flexible \implies
- 2 M is not **c**-flexible \implies

FLEXIBILITY IN DIMENSION 6

Relations:

$$\begin{aligned}\hat{c}_3 &= c_4^{(1)} + 1 - b_1 - b_2 + b_3, \\ \hat{c}_3 &= h_3 - h_5, \\ \check{c}_4^{(1)} &= \hat{c}_3 - h_4 + b_2, \\ \check{c}_5^{(1)} &= c_4^{(1)}, \\ \check{c}_5^{(2)} &= b_1 + b_2 - h_5 - 1.\end{aligned}$$

Results:

- 1 M is **c**-flexible $\implies M$ is **f**-flexible and **h**-flexible.
- 2 M is not **c**-flexible \implies

FLEXIBILITY IN DIMENSION 6

Relations:

$$\begin{aligned}\hat{c}_3 &= c_4^{(1)} + 1 - b_1 - b_2 + b_3, \\ \hat{c}_3 &= h_3 - h_5, \\ \check{c}_4^{(1)} &= \hat{c}_3 - h_4 + b_2, \\ \check{c}_5^{(1)} &= c_4^{(1)}, \\ \check{c}_5^{(2)} &= b_1 + b_2 - h_5 - 1.\end{aligned}$$

Results:

- 1 M is **c**-flexible $\implies M$ is **f**-flexible and **h**-flexible.
- 2 M is not **c**-flexible $\implies M$ is **f**-flexible if and only if it is **h**-flexible.

FLEXIBILITY IN DIMENSION 6

Relations:

$$\begin{aligned}\hat{c}_3 &= c_4^{(1)} + 1 - b_1 - b_2 + b_3, \\ \hat{c}_3 &= h_3 - h_5, \\ \check{c}_4^{(1)} &= \hat{c}_3 - h_4 + b_2, \\ \check{c}_5^{(1)} &= c_4^{(1)}, \\ \check{c}_5^{(2)} &= b_1 + b_2 - h_5 - 1.\end{aligned}$$

Results:

- 1 M is **c**-flexible $\implies M$ is **f**-flexible and **h**-flexible.
- 2 M is not **c**-flexible $\implies M$ is **f**-flexible if and only if it is **h**-flexible.

Examples of **h**-flexible nilmanifolds in dimension 6 can be found in [IRTU 01].

We classify the 6-dimensional **c**-flexible nilmanifolds.

FLEXIBILITY IN DIMENSION ≥ 8

Let (M^{2n}, ω) be a closed symplectic manifold.

- 1 M is **c**-flexible $\implies M$ is **f**-flexible or **h**-flexible.
- 2 $\exists M$ **f-flexible** for any dimension $2n$.

Example in dimension 8: a solvmanifold that is **c**-flexible, **f**-flexible and **h**-flexible.